

cAMP/PKA pathway activation in human mesenchymal stem cells *in vitro* results in robust bone formation *in vivo*

Ramakrishnaiah Siddappa*, Anton Martens†, Joyce Doorn*, Anouk Leusink*, Cristina Olivo†, Ruud Licht*, Linda van Rijn*, Claudia Gaspar‡, Riccardo Fodde‡, Frank Janssen*, Clemens van Blitterswijk*, and Jan de Boer*[§]

*Department of Tissue Regeneration, Institute for Biomedical Technology, University of Twente, 7500 AE, Enschede, The Netherlands; †Department of Immunology, University Medical Center Utrecht, 3584 EA, Utrecht, The Netherlands; and ‡Department of Pathology, Josephine Nefkens Institute, Erasmus Medical Center, 3000 DR, Rotterdam, The Netherlands

Edited by Robert Langer, Massachusetts Institute of Technology, Cambridge, MA, and approved March 17, 2008 (received for review November 29, 2007)

Tissue engineering of large bone defects is approached through implantation of autologous osteogenic cells, generally referred to as multipotent stromal cells or mesenchymal stem cells (MSCs). Animal-derived MSCs successfully bridge large bone defects, but models for ectopic bone formation as well as recent clinical trials demonstrate that bone formation by human MSCs (hMSCs) is inadequate. The expansion phase presents an attractive window to direct hMSCs by pharmacological manipulation, even though no profound effect on bone formation *in vivo* has been described so far using this approach. We report that activation of protein kinase A elicits an immediate response through induction of genes such as *ID2* and *FosB*, followed by sustained secretion of bone-related cytokines such as BMP-2, IGF-1, and IL-11. As a consequence, PKA activation results in robust *in vivo* bone formation by hMSCs derived from orthopedic patients.

bone tissue engineering | osteogenesis | PKA signaling

The ability of human mesenchymal stem cells (hMSCs) to differentiate into adipogenic, chondrogenic, osteogenic (1), and myogenic (2) lineages has generated a great deal of potential clinical use in regenerative medicine and tissue engineering in the past decade. Concomitantly, hMSCs are increasingly used as a cell biological model system to investigate molecular mechanisms governing signal transduction (3), differentiation (4–6), cell fate decision (7), and senescence (8, 9) because the step from basic research to clinical application is relatively short. Although hMSCs are multipotent and form mineralized bone tissue *in vivo*, their clinical application is still limited. Using a bone tissue engineering approach, large bone defects can be repaired in animal models by using animal-derived MSCs. For instance, Petite *et al.* (10) demonstrated bone union in a metatarsal defect in sheep. In contrast, bone formation by hMSCs is substantially less robust, and repair of critical size bone defects has not been achieved by hMSCs so far (11).

Predifferentiation of hMSCs into the osteogenic lineage *in vitro* during the expansion phase before implantation offers an opportunity to improve their *in vivo* performance. In previous studies, dexamethasone (dex) and vitamin D3 were used to promote hMSC differentiation *in vitro* (12). More recent studies include the MAPK pathway (13), Rho kinase (7), Wnt (5), Notch (14), and receptor tyrosine kinases (3). So far, no positive correlation between hMSC osteogenesis *in vitro* and bone formation *in vivo* has been reported. Although many reports describe the positive effect of compounds on osteogenesis *in vitro*, there are, to the best of our knowledge, no convincing reports where *in vitro* manipulation of clinically applicable hMSCs significantly augments bone formation *in vivo*. Either no *in vivo* experiments were performed or the effect on bone formation is marginal and sometimes even negative (15, 16). In other studies, transgenic immortalized hMSC derivative was used (3, 9).

Protein kinase A (PKA) signaling plays a prominent but ambiguous role in mesenchymal cell fate decision, which depends on the molecular and developmental context in which the PKA signal is

presented (17, 18). Relatively little is known about the role of PKA in osteogenic differentiation of hMSCs, but it is anticipated by the anabolic effect of certain PKA-activating hormones on bone mineral density. Intermittent administration of parathyroid hormone increases trabecular and cancellous bone formation in ovariectomized mice, although continuous administration results in net bone loss (19). The effect of PKA activation on osteogenesis has been studied in different cell types with compounds that directly or indirectly activate PKA, although the results are contentious. The most direct evidence on a role of PKA in osteogenic differentiation is from studies in calcifying vascular cells (20). Here activation of the PKA pathway with *N*⁶,2'-*O*-dibutyryladenosine-3',5'-cyclic monophosphate (db-cAMP) stimulated the expression of osteogenic marker genes and *in vitro* mineralization, suggesting that the PKA pathway promotes vascular calcification by enhancing osteogenic differentiation of calcifying vascular cells. Furthermore, the PKA activator forskolin increased bone nodule formation at low concentrations but inhibited it at higher concentrations (21). A recent study shows that parathyroid hormone-related peptide (PTHrP) inhibits *CBFA1* expression through the PKA pathway (22), and it has been reported that PKA activation enhances adipogenic differentiation of hMSCs (18). In this article we describe that PKA-activated hMSCs demonstrate enhanced *in vitro* osteogenesis and *in vivo* bone formation, which opens a promising window of opportunity to further improve bone tissue engineering protocols.

Results

cAMP/PKA Signaling Induces Osteogenesis in hMSCs. To assess the effect of PKA activation on osteogenesis in hMSCs, we exposed a panel of hMSCs, isolated from the bone marrow of 14 patients undergoing orthopedic surgery ranging 31–82 years of age [supporting information (SI) Table S1], to the PKA activator db-cAMP. Both dex and db-cAMP consistently enhanced the expression of the early osteogenic marker alkaline phosphatase (ALP), ranging from a 1.8-fold increase to a 5.3-fold increase compared with untreated controls (see Fig. 1*a* and Fig. S1*a*). Donor variation was observed both in the basal level of ALP expression, as reported by us previously (23), and in their response to both agents (Fig. S1*a*). db-cAMP-induced ALP expression did not depend on the presence of dex, although coexposure resulted in an additive and sometimes

Author contributions: R.S., C.v.B., and J.d.B. designed research; R.S., J.D., R.L., L.v.R., and F.J. performed research; A.M., C.O., R.L., C.G., and R.F. contributed new reagents/analytic tools; R.S. and A.L. analyzed data; and R.S., C.v.B., and J.d.B. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Freely available online through the PNAS open access option.

[§]To whom correspondence should be addressed. E-mail: j.deboer@tnw.utwente.nl.

This article contains supporting information online at www.pnas.org/cgi/content/full/0711190105/DCSupplemental.

© 2008 by The National Academy of Sciences of the USA




Fig. 1. PKA activation induces *in vitro* osteogenesis of hMSCs. (a) Box plot showing the average percentage of ALP-positive cells from 14 donors in basic medium (Con), osteogenic medium (Dex), basic medium with 1 mM db-cAMP (cAMP), or osteogenic medium supplemented with 1 mM db-cAMP (Dex+cAMP). The data were analyzed by using two-way ANOVA followed by Dunnett's multiple-comparison test. Statistical significance is denoted compared with the control group. (b) hMSCs were grown in either mineralization medium (dex) or mineralization medium to which 1 mM db-cAMP was added during the first 3, 5, 10, 15, 25, or full 30 days after which calcium deposition was measured and expressed as micrograms of calcium per milliliter of sample. The data were analyzed by using one-way ANOVA followed by Dunnett's multiple-comparison test. (c) H89, a PKA inhibitor, reverses the db-cAMP-induced ALP expression. hMSCs were preincubated with H89 for 10–15 h and then cotreated with db-cAMP or cholera toxin (CTX) for 4 days. The data were analyzed by using one-way ANOVA followed by Tukey's multiple-comparison test. (d) Addition of db-cAMP to hMSCs for 6 h resulted in increased phosphorylation of transcription factor CREB, which could be inhibited by coinubation with H89. *, $P < 0.05$; **, $P < 0.01$.

synergistic effect on ALP expression, resulting in up to 60% of ALP-positive cells in some donors. Consistent with a positive effect on ALP expression, calcium deposition was enhanced when db-cAMP was administered to hMSCs in mineralization medium (Fig. 1b). Optimal stimulation of mineralization was observed when hMSCs were exposed to db-cAMP during the first 5 days of osteogenic culture, resulting in a 3-fold increase in mineralization compared with treatment with dex alone. Incubation for >15 days resulted in a significant inhibition of mineralization. The positive effect of PKA signaling on osteogenesis was further confirmed by quantitative PCR (qPCR) on a panel of osteogenic marker genes during a 15-day osteogenic time course. db-cAMP had a significant positive effect on *ALP* and *COL1A1* expression from day 3 until day 15 (Fig. S2). Dex and db-cAMP together induced the expression of the transcription factor *CBFA1* on days 5 and 15. In contrast, *OPN* expression was induced only at day 10 by dex or dex and db-cAMP and appears to be a dex-mediated event. *BGLAP*, *ON*, and *S100A4* showed no consistent difference in expression profile during the course of osteogenic differentiation.

To demonstrate that db-cAMP mediates its effect through activation of PKA, we exposed cells to two other upstream PKA

activators, cholera toxin and forskolin, and confirmed that also these compounds stimulate ALP expression in hMSCs (Fig. 1c and Fig. S1b). When hMSCs were exposed to db-cAMP or cholera toxin in the presence of H89, an inhibitor of PKA activity, PKA-induced ALP was reduced to basal level. Interestingly, H89 did not significantly affect dex-induced ALP expression, showing that dex and db-cAMP stimulate ALP expression through discrete molecular mechanisms (Fig. 1c). One of the direct target proteins of PKA is cAMP response element-binding protein (CREB), and indeed we detected phosphorylated CREB in hMSCs upon treatment with db-cAMP (Fig. 1d). Evidently, treatment of hMSCs with db-cAMP leads to PKA activation, which stimulates *in vitro* osteogenic differentiation of hMSCs.

cAMP/PKA Signaling Enhances *in Vivo* Bone Formation by hMSCs. To evaluate the effect of db-cAMP on *in vivo* bone formation we used the ectopic bone formation model in immune-deficient mice, which is widely used to assess the bone-forming capacity of hMSCs (3, 9, 11, 24). Using this model, we typically observe a bone/ceramic surface ratio of 15–20% for goat MSCs, as shown in Fig. 2a. This represents $\approx 50\%$ of the available pore area, and all of the pores

to inhibit cell proliferation (25). Despite the negative on proliferation, db-cAMP-treated hMSCs produced three times more bone (15% bone/ceramic surface ratio) than untreated control cells (5%), which is equal to the amount of bone deposited by goat MSCs (Fig. 2a). Histological examination of the explanted grafts showed that mature bone was formed in which osteocytes were embedded in mineralizing extracellular matrix (Fig. 2f) and mineralizing osteoblasts could be detected. Polarizing light microscopy of the deposited matrix showed areas of lamellar bone, which indicates that bone tissue has been remodeled by osteoclasts and osteoblasts (Fig. S4). Moreover, marrow-like tissue was observed in the vicinity of the tissue-engineered bone, which further indicates the functionality of the bone tissue (Fig. 2g) (24).

Autocrine and Paracrine Induction of Bone Formation by cAMP/PKA-Activated hMSCs. Based on the robust effect of db-cAMP on bone formation *in vivo*, which is in striking contrast to the effect of dex, we anticipated that PKA activation results in the expression of a unique set of target genes ultimately resulting in osteogenic differentiation *in vitro* and bone formation *in vivo*. To get more insight into the molecular mechanism behind db-cAMP-induced bone formation, we compared the gene expression profile of untreated cells with that of hMSCs treated with db-cAMP for 6 h using DNA microarray technology. Among the 62 genes up-regulated six times or more by db-cAMP (Table S2), a number of genes were previously associated with cAMP/PKA signaling. Six genes were identified in chip-on-chip assays for CREB-binding promoters (*CREM*, *RCC1*, *GP-1*, *FLRT3*, *GPRC5A*, and *EVX-1*; see <http://natural.salk.edu/CREB/>) whereas other genes have been implicated in cAMP/PKA signaling, e.g., *ID-2* (26), *BMP-2* (27), *IGF-1* (28), and *IL-11* (29). To show that the genes are directly activated by PKA signaling, we demonstrated that db-cAMP-mediated *ID-2* expression is insensitive to cycloheximide, an inhibitor of translation (Fig. 3a). Gene Ontology enrichment of the data set further showed a remarkably high number of transcription factors (nine of the top 62 up-regulated genes). Three transcription factors are implicated in osteogenesis [*c-Fos*, *FosB*, and *ID-2* (30)], but others were previously not implicated in osteogenesis, such as *BCL3*, *EVX1*, *CDX2*, *DUX1*, *DAT1*, and *ID-4*. We did not observe the classical osteogenic transcription factors such as *CBFA1* (31) and *OSX/SP7* (32) in our microarray data set. We also analyzed the gene expression profile of hMSCs at the time of implantation. Gene Ontology enrichment of genes up-regulated five times or more in hMSCs treated for 7 days with db-cAMP shows that, at this point in time, hMSCs express many secreted proteins, mostly growth factors and cytokines (Table S3). Most striking is the 46-fold up-regulation of *BMP-2*, which suggests that autocrine or paracrine BMP signaling is involved in the osteogenic effect of db-cAMP. To investigate this, we analyzed the expression of *ID-1*, a typical BMP target gene, by qPCR during db-cAMP-driven *in vitro* differentiation. When hMSCs were exposed to db-cAMP and dex, *ID-1* expression was significantly higher on day 5 compared with db-cAMP alone. Dex alone did not affect *ID-1* expression compared with untreated cells (Fig. 3b). Other cytokines that have been implicated in osteogenesis and bone formation (*IGF-1*, *IL-8*, and *IL-11*) (33) ranked among the most highly regulated genes (19-, 47-, and 18-fold up-regulated, respectively; see Table S3). We confirmed the secretion of these cytokines by db-cAMP-treated hMSCs using ELISA (Fig. 3c). In this case, the induction is even more drastic. Another interesting set of cytokines up-regulated in the 7-day treatment group are proteins interfering with vasculogenesis such as angiopoietin-like 2, angiopoietin-like 4, and placental growth factor. Based on the putative involvement of BMP and other secreted proteins, we propose a model for db-cAMP-induced bone formation in which paracrine signaling through hMSC-secreted cytokines and growth factors stimulates bone formation *in vivo* (Fig. 4).




Fig. 3. db-cAMP-induced gene and protein expression. (a) hMSCs were treated with cycloheximide for 1 h and then coincubated with db-cAMP for 6 more hours. Expression of BMP target genes *ID-1* and *ID-2* was analyzed compared to cycloheximide-treated cells. The data were analyzed by using Student's *t* test. (b) hMSCs were grown in basic medium, basic medium supplemented with 1 mM db-cAMP (cAMP), osteogenic medium (Dex), or osteogenic medium supplemented with 1 mM db-cAMP (Dex+cAMP). Expression of *ID-1* was analyzed by qPCR and is expressed as fold induction compared with cells grown in basic medium. The data were analyzed by using two-way ANOVA. Statistical differences are denoted compared with cells grown in basic medium. (c and d) db-cAMP induces secretion of proosteogenic cytokines and growth factors. hMSCs were treated with db-cAMP for 4 days, the supernatant was collected, and IGF-1 (c), IL-8, and IL-11 (d) expression in the medium was measured by ELISA. The data were analyzed by using Student's *t* test. **, $P < 0.01$.

Discussion

The osteogenic potential of hMSCs has been long recognized and provides the exciting potential to treat patients with bone defects beyond spontaneous healing (1, 34). Successful bone formation has been reported for animal models, but so far bone formation by hMSCs is limited and recent clinical trials have demonstrated that *in vivo* bone formed by the implanted hMSCs is insufficient (11). Bone tissue engineering may be augmented through systemic administration of osteogenic agents or by scaffold-mediated delivery of biologically active reagents (35). Manipulation of hMSCs during *in vitro* expansion is an interesting alternative because, in this approach, the agents used *in vitro* can be cleared from the graft before implantation with minimal regulatory and safety implications. As such, large libraries of small molecules can be screened and used for their desired biological effect (36). The lack of irrefutable data demonstrating that predifferentiation of hMSCs *in vitro* results in enhanced bone formation *in vivo* led us to investigate whether db-cAMP treatment enhances *in vivo* performance of hMSCs. Our data describe the osteogenic effect of PKA activation in hMSCs, and we demonstrate that *in vitro* manipulation of hMSCs with the small molecule db-cAMP results in robust osteogenic differentiation *in vitro* and bone formation *in vivo*. The mechanism of action seems to be different from that used by dex and may involve paracrine/autocrine signaling by BMPs and other osteogenic cytokines. However, other mechanisms such as improved survival of db-cAMP-treated cells or enhanced angiogenesis cannot be ex-




Fig. 4. Model for autocrine/paracrine induction of osteogenesis in hMSCs by PKA signaling. db-cAMP induces direct expression of BMP target genes such as *ID-2* and *ID-4* via CREB resulting in cell-autonomous stimulation of osteogenesis whereas expression of BMP-2, proosteogenic cytokines, and growth factors results in paracrine induction of bone formation.

cluded, which warrants further research to confirm our model (Fig. 4).

We evaluated the potential application of db-cAMP-treated hMSCs in bone tissue engineering by testing its effect on *in vivo* bone formation using different tissue engineering strategies currently in use: standard tissue engineering, peroperative seeding, and bioreactor-based bone tissue engineering. In the peroperative seeding strategy, hMSCs are poli-clinically isolated from a bone marrow biopsy or alternative sources such as fat tissue (37) and expanded. On the day of the surgery, cells are seeded onto the scaffold material and implanted in the defect (38). In addition, automated bioreactor systems may play a role in future bone tissue engineering because a clinical research team can harvest a biopsy and expand and differentiate cell-seeded grafts in routine clinical laboratories up to the time of implantation. Prototypes of bioreactors have been reported by us and others (39, 40). In our hands, bioreactor-mediated bone tissue engineering is the most efficient method. We are trying to enhance bone formation even further by improving the proliferation phase of MSCs before db-cAMP treatment and by finding the most optimal balance between induction of osteogenesis and cell proliferation. With further improvement on the proliferation and differentiation scheme, we hope to consistently achieve a 15–20% bone/ceramic surface ratio.

The data presented in this article describe for the first time that bone formation by hMSCs can be significantly augmented through manipulation of the signal transduction milieu *in vitro* by using a simple compound like db-cAMP. This work once again demonstrates the enormous plasticity of hMSCs and strongly encourages further efforts to engineer the gene expression profile of hMSCs for optimal clinical application.

Materials and Methods

Isolation, Culture, and ALP Analysis of hMSCs. Bone marrow aspirates were obtained from donors with written informed consent, and hMSCs were isolated

and proliferated as described previously (41). Briefly, aspirates were resuspended by using 20-gauge needles, plated at a density of 5×10^5 cells per square centimeter and cultured in hMSC proliferation medium containing α -MEM (Life Technologies), 10% FBS (Cambrex), 0.2 mM ascorbic acid (Asap; Life Technologies), 2 mM L-glutamine (Life Technologies), 100 units/ml penicillin (Life Technologies), 10 μ g/ml streptomycin (Life Technologies), and 1 ng/ml basic FGF (Instruchemie). Cells were grown at 37°C in a humid atmosphere with 5% CO₂. Medium was refreshed twice a week, and cells were used for further subculturing or cryopreservation. hMSC basic medium/control medium was composed of hMSC proliferative medium without basic FGF, hMSC osteogenic medium was composed of hMSC basic medium supplemented with 10^{-8} M dex (Sigma), and hMSC mineralization medium was composed of basic medium supplemented with 10^{-8} M dex and 0.01 M β -glycerophosphate (Sigma). Donor information is provided in Table S1.

Osteogenic Assay. To determine whether PKA activation elicits an osteogenic response in hMSCs, we exposed them to 1 mM db-cAMP (Sigma) with or without dex for 4 days and analyzed the expression of the osteogenic marker ALP by flow cytometry. hMSCs were seeded at 5,000 cells per square centimeter and allowed to attach for 10–15 h in basic medium, then cells were incubated with 10^{-8} M dex and 1 mM db-cAMP for the denoted time periods. Each experiment was performed in triplicate with a negative control (cells grown in basic medium) and a positive control (cells grown in the presence of dex) and one or more experimental conditions. At the end of the culture period, the cells were trypsinized and incubated for 30 min in block buffer [PBS with 5% BSA (Sigma) and 0.05% NaN₃], then incubated with primary antibody (anti-ALP, B4-78; Developmental Studies Hybridoma Bank, University of Iowa) diluted in wash buffer (PBS with 1% BSA and 0.05% NaN₃) for 30 min or with isotype control antibodies. Cells were then washed three times with wash buffer and incubated with secondary antibody (rabbit anti-mouse IgG phycoerythrin; DAKO) for 30 min. Cells were washed three times and suspended in 250 μ l of wash buffer with 10 μ l of Viaprobe (Pharmin-gen) for live/dead cell staining, and only live cells were used for further analysis. The percentage of ALP-positive cells was calculated on a FACSCalibur (Becton Dickinson Immunocytometry Systems). Mineralization was measured quantitatively by using a calcium assay kit (Sigma Diagnostics; 587A) according to the manufacturer's protocol after culturing the cells for the denoted time period with the compounds mentioned.

Gene Expression Analysis by qPCR and Microarray. The effect of db-cAMP on expression of osteogenic marker genes was analyzed by seeding hMSCs at 5,000 cells per square centimeter in T75 flasks supplemented in various medium compositions for 3, 5, 10, and 15 days. To analyze the direct induction of BMP target genes by PKA activation, hMSCs were seeded at 5,000 cells per square centimeter and supplemented with 1 mM db-cAMP for 6 h with or without 10 μ M cycloheximide (Sigma). RNA was isolated by using an RNeasy mini kit (Qiagen), and qPCR was performed by using SYBR green (Invitrogen) on a Light Cycler (Roche). Data were analyzed by using Light Cycler software version 3.5.3, using the fit point method by setting the noise band to the exponential phase of the reaction to exclude background fluorescence. Expression of osteogenic marker genes was calculated relative to 18s rRNA levels by the comparative Δ CT method (42). To study the genome-wide effect of db-cAMP, hMSCs were grown in either basic medium or basic medium supplemented with 1 mM db-cAMP for 6 h or 7 days. RNA was isolated by using an RNeasy midi kit (Qiagen), and 8 μ g of total RNA was used for probe labeling according to the manufacturer's protocol (Affymetrix). The probe quality was verified by using lab-on-chip technology (Agilent Technologies), and samples were hybridized to Human Genome Focus arrays according to the manufacturer's protocol (Affymetrix). Data analysis was performed by using Affymetrix GENECHIP 4.0 software. CREB (Calbiochem), phosphorylated CREB (R & D Systems), and β -actin (R & D Systems) antibodies were used to detect respective proteins by Western blotting on cell lysates obtained from hMSCs treated with various supplements. IGF-1, IL-8, and IL-11 secretion upon db-cAMP treatment was measured in the cell supernatant by ELISA (R & D Systems) according to the manufacturer's protocol.

In Vivo Evaluation Studies. To evaluate the effect of PKA activation on ectopic bone formation by hMSCs, we used three tissue engineering protocols.

Standard tissue engineering approach. As scaffold for all bone tissue engineering methods, we used porous biphasic calcium phosphate (BCP) ceramic granules of \approx 2–3 mm, prepared and sintered at 1,150°C as described previously (43). The cells were cultured for 7 days in basic medium, in osteogenic medium, or in osteogenic medium supplemented with 1 mM db-cAMP during the last 4 days. Goat MSCs were isolated and expanded as described previously (40).

Peroperative seeding approach. To investigate the performance of db-cAMP-treated hMSCs in the peroperative seeding approach, we expanded hMSCs from eight donors and changed the medium to basic medium or medium containing 1 mM db-cAMP 4 days before implantation. On the day of the operation, we

trypsinized the cells and allowed them to attach to porous BCP scaffolds for 4 h as described above.

Bioreactor-based tissue engineering. hMSCs were seeded onto porous ceramics as described above, transferred into a bioreactor, and cultured for 4 days in basic medium. A direct perfusion flow bioreactor was used as described previously (40). Briefly, the bioreactor comprised an inner and outer housing, which are configured as coaxially disposed, nested cylinders. The bioreactor system consisted of a bioreactor, a sterile fluid pathway (made of γ -sterilized PVC tubing with low gas permeability) that includes a medium supply vessel, a pump, an oxygenator, and a waste vessel. The fluid pathway contained a temperature sensor and two dissolved oxygen sensors, which are placed at the medium inlet and outlet of the bioreactor. The whole bioreactor system was placed in a temperature-controlled unit at 37°C. The incubation units were equipped with controlled oxygen and carbon dioxide supplying systems. The gas environment in the chamber is kept at a constant level of 21% O₂ and 5% CO₂, and medium is pumped through the gas-permeable tube. This system enables a medium flow over and through the cell-seeded biomaterials with constant pH and a constant oxygen concentration. During the last 4 days, the cells in the bioreactor were or were not supplemented with 1 mM db-cAMP perfused through the medium.

At the end of the culture period, the tissue-engineered constructs were implanted s.c. in immune-deficient mice for 6 weeks. In each *in vivo* experiment, 10 nude male mice (Hsd-cpb:NMRI-nu; Harlan) were anesthetized by intramuscular injection of 0.05 ml of anesthetic (1.75 ml of 100 μ g/ml ketamine, 1.5 ml of 20 mg/ml xylazine, and 0.5 ml of 0.5 mg/ml atropine). Four s.c. pockets were made, and each pocket was implanted with three particles of each condition. The incisions were closed with a vicryl 5-0 suture, and the tissue-engineered constructs were left for 6 weeks. All experiments were approved by the local Animal Experimental Committee.

Histology and Histomorphometry. After 6 weeks, the mice were killed by using CO₂ and samples were explanted, fixed in 1.5% glutaraldehyde (Merck) in 0.14 M cacodylic acid (Fluka) buffer (pH 7.3), dehydrated, and embedded in methyl methacrylate (Sigma) for sectioning. Approximately 10- μ m-thick, undecalcified sections were processed on a histological diamond saw (Leica saw microtome cutting system). The sections were stained with basic fuchsin and methylene blue

to visualize new bone formation. The newly formed mineralized bone stains red with basic fuchsin, all other cellular tissues stain light blue with methylene blue, and the ceramic material remains black and unstained by both the dyes. Histological sections were qualitatively analyzed by using a light microscope (Leica), and each histological section was scored either positive or negative for bone formation. Quantitative histomorphometry was performed as described previously (44). Briefly, high-resolution digital photographs (300 dpi) were made from four randomly selected sections from each tissue-engineered graft. Before histomorphometrical analysis, bone and material were pseudocolored green and red, respectively, by using Photoshop CS2 (Adobe Systems). Image analysis was performed by using a PC-based system with KS400 software (version 3, Zeiss). A custom-made macro was used to measure bone/ceramic surface ratios.

Cell distribution on the scaffold and matrix formation were qualitatively visualized with methylene blue staining. The tissue-engineered constructs were immersed in 1% methylene blue solution for 1 min and washed three times with demineralized water to remove unbound stain. The BCP particles remain unstained while cells stain blue when examined by light microscopy. To assess the effect of db-cAMP on proliferation, cells were seeded in basic medium at 5,000 cells per square centimeter either with or without 1 mM db-cAMP for 5 days. Next, the medium was removed and 2 ml of a 10% Alamar blue (Biosource) solution was added and incubated for 4 h. From this, 0.2 ml was transferred to a 96-well plate and measured on a spectrophotometer (PerkinElmer) at 545 nm.

ACKNOWLEDGMENTS. We thank Dr. L. Creemers and Dr. W. Dhert (University Medical Center Utrecht) and Dr. A. Renard (Medisch Spectrum Twente, Enschede, The Netherlands) for kindly providing us with bone marrow aspirates and Dr. M. Karperien for critically reading the manuscript. We also thank Sanne Both, Lotus Sterk, Hugo Alves, Hugo Fernandes, Remi Tibben, and Roland Heerkens for technical support and Huipin Yuan (University of Twente) for supply of ceramic materials. The work was supported by grants from The Netherlands Ministry of Economic Affairs (SenterNovem; to R.S., R.L., and J.d.B.), The Netherlands Organization for Scientific Research (NWO/Vici 016.036.636 to R.F.), Grant 03038 from the Besluit Subsidies Investeren Kennisinfrastructuur program of the Dutch Government (to R.F.), the European Union FP6 Migrating Cancer Stem Cells Program (R.F.), the Dutch Program for Tissue Engineering (C.O. and A.M.), and the Italian Association for Research on Cancer (C.G.).

- Pittenger MF, et al. (1999) Multilineage potential of adult human mesenchymal stem cells. *Science* 284:143–147.
- Dezawa M, et al. (2005) Bone marrow stromal cells generate muscle cells and repair muscle degeneration. *Science* 309:314–317.
- Kratchmarova I, et al. (2005) Mechanism of divergent growth factor effects in mesenchymal stem cell differentiation. *Science* 308:1472–1477.
- Boland GM, Perkins G, Hall DJ, Tuan RS (2004) Wnt 3a promotes proliferation and suppresses osteogenic differentiation of adult human mesenchymal stem cells. *J Cell Biochem* 93:1210–1230.
- de Boer J, et al. (2004) Wnt signaling inhibits osteogenic differentiation of human mesenchymal stem cells. *Bone* 34:818–826.
- de Boer J, Wang HJ, van Blitterswijk C (2004) Effects of Wnt signaling on proliferation and differentiation of human mesenchymal stem cells. *Tissue Eng* 10:393–401.
- McBeath R, et al. (2004) Cell shape, cytoskeletal tension, and RhoA regulate stem cell lineage commitment. *Dev Cell* 6:483–495.
- Shi S, et al. (2002) Bone formation by human postnatal bone marrow stromal stem cells is enhanced by telomerase expression. *Nat Biotechnol* 20:587–591.
- Simonsen JL, et al. (2002) Telomerase expression extends the proliferative life-span and maintains the osteogenic potential of human bone marrow stromal cells. *Nat Biotechnol* 20:592–596.
- Petite H, et al. (2000) Tissue-engineered bone regeneration. *Nat Biotechnol* 18:959–963.
- Meijer GJ, de Bruijn JD, Koole R, van Blitterswijk CA (2007) Cell-based bone tissue engineering. *PLoS Med* 4:e9.
- Siddappa R, et al. (2007) The response of human mesenchymal stem cells to osteogenic signals and its impact on bone tissue engineering. *Curr Stem Cell Res Ther* 2:209–220.
- Jaiswal RK, et al. (2000) Adult human mesenchymal stem cell differentiation to the osteogenic or adipogenic lineage is regulated by mitogen-activated protein kinase. *J Biol Chem* 275:9645–9652.
- Tezuka K, et al. (2002) Stimulation of osteoblastic cell differentiation by Notch. *J Bone Miner Res* 17:231–239.
- de Boer J, et al. (2006) Inhibition of histone acetylation as a tool in bone tissue engineering. *Tissue Eng* 12:2927–2937.
- de Kok IJ, et al. (2006) Effect of vitamin D pretreatment of human mesenchymal stem cells on ectopic bone formation. *J Oral Maxillofac Surg* 64:103–109.
- Yang X, et al. (2004) ATF4 is a substrate of RSK2 and an essential regulator of osteoblast biology; implication for Coffin-Lowry syndrome. *Cell* 117:387–398.
- Zhao Y, Ding S (2007) A high-throughput siRNA library screen identifies osteogenic suppressors in human mesenchymal stem cells. *Proc Natl Acad Sci USA* 104:9673–9678.
- Jilka RL (2007) Molecular and cellular mechanisms of the anabolic effect of intermittent PTH. *Bone* 40:1434–1446.
- Tintut Y, et al. (1998) cAMP stimulates osteoblast-like differentiation of calcifying vascular cells: Potential signaling pathway for vascular calcification. *J Biol Chem* 273:7547–7553.
- Turksen K, Grigoriadis AE, Heersche JN, Aubin JE (1990) Forskolin has biphasic effects on osteoprogenitor cell differentiation in vitro. *J Cell Physiol* 142:61–69.
- Li TF, et al. (2004) Parathyroid hormone-related peptide (PTHrP) inhibits Runx2 expression through the PKA signaling pathway. *Exp Cell Res* 299:128–136.
- Siddappa R, Licht R, van Blitterswijk C, de Boer J (2007) Donor variation and loss of multipotency during *in vitro* expansion of human mesenchymal stem cells for bone tissue engineering. *J Orthop Res* 25:1029–1041.
- Sacchetti B, et al. (2007) Self-renewing osteoprogenitors in bone marrow sinusoids can organize a hematopoietic microenvironment. *Cell* 131:324–336.
- Del Sal G, et al. (1995) Gas1-induced growth suppression requires a transactivation-induced p53 function. *Mol Cell Biol* 15:7152–7160.
- Kurabayashi M, Dutta S, Jeyaseelan R, Kedes L (1995) Doxorubicin-induced Id2A gene transcription is targeted at an activating transcription factor/cyclic AMP response element motif through novel mechanisms involving protein kinases distinct from protein kinase C and protein kinase A. *Mol Cell Biol* 15:6386–6397.
- Ionescu AM, et al. (2004) CREB cooperates with BMP-stimulated Smad signaling to enhance transcription of the Smad6 promoter. *J Cell Physiol* 198:428–440.
- Umayahara Y, et al. (1997) CCAAT/enhancer-binding protein delta activates insulin-like growth factor-I gene transcription in osteoblasts: Identification of a novel cyclic AMP signaling pathway in bone. *J Biol Chem* 272:31793–31800.
- Kim GS, et al. (1997) Involvement of different second messengers in parathyroid hormone- and interleukin-1-induced interleukin-6 and interleukin-11 production in human bone marrow stromal cells. *J Bone Miner Res* 12:896–902.
- Karsenty G, Wagner EF (2002) Reaching a genetic and molecular understanding of skeletal development. *Dev Cell* 2:389–406.
- Ducy P, et al. (1997) Osf2/Cbfa1: A transcriptional activator of osteoblast differentiation. *Cell* 89:747–754.
- Nakashima K, et al. (2002) The novel zinc finger-containing transcription factor osterix is required for osteoblast differentiation and bone formation. *Cell* 108:17–29.
- Hughes FJ, Turner RT, Belibasakis G, Martuscelli G (2006) Effects of growth factors and cytokines on osteoblast differentiation. *Periodontol* 2000 41:48–72.
- Bianco P, Robey PG (2001) Stem cells in tissue engineering. *Nature* 414:118–121.
- Lutolf MP, Hubbell JA (2005) Synthetic biomaterials as instructive extracellular microenvironments for morphogenesis in tissue engineering. *Nat Biotechnol* 23:47–55.
- Ding S, Schultz PG (2004) A role for chemistry in stem cell biology. *Nat Biotechnol* 22:833–840.
- Cowan CM, et al. (2004) Adipose-derived adult stromal cells heal critical-size mouse calvarial defects. *Nat Biotechnol* 22:560–567.
- Krutz MC, et al. (2004) Optimization of bone tissue engineering in goats: A peroperative seeding method using cryopreserved cells and localized bone formation in calcium phosphate scaffolds. *Transplantation* 77:359–365.
- Braccini A, et al. (2005) Three-dimensional perfusion culture of human bone marrow cells and generation of osteoinductive grafts. *Stem Cells* 23:1066–1072.
- Janssen FW, Oostra J, Oorschot A, van Blitterswijk CA (2006) A perfusion bioreactor system capable of producing clinically relevant volumes of tissue-engineered bone: *In vivo* bone formation showing proof of concept. *Biomaterials* 27:315–323.
- Both SK, et al. (2007) A rapid and efficient method for expansion of human mesenchymal stem cells. *Tissue Eng* 13:3–9.
- Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 25:402–408.
- Yuan H, et al. (2002) A comparison of the osteoinductive potential of two calcium phosphate ceramics implanted intramuscularly in goats. *J Mater Sci Mater Med* 13:1271–1275.
- Habibovic P, et al. (2008) Osteoconduction and osteoinduction of low-temperature 3D printed bioceramic implants. *Biomaterials* 29:944–953.